abcam

ab92513

Anti-ERG antibody [EPR3864]

Anti-ERG antibody [EPR3864] (ab92513) is a rabbit monoclonal antibody detecting ERG in **Western Blot, Flow Cytometry (Intra), IHC-P, ICC/IF**. Suitable for **Human, Mouse, Rat**.

- Clone EPR3864 is the most cited clone to ERG
- Biophysical QC for unrivalled batch-batch consistency
- Over 230 publications
- Trusted since 2010

Mouse version of this clone also available as ab214341. Same clone, same staining pattern, different species antibody backbone.
Learn more about chimeric antibodies

Recombinant | RabMAb | Advanced Validation

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR3864
Purification technique	Affinity purification Protein A
Specificity	This antibody also detects Fli-1.
Dissociation constant	8.9×10^{-10} M
Concentration	0.903 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Mouse
Dilution info	1/1000
Notes	**For unpurified, use 1/100 - 1/250.** Perform heat-mediated antigen retrieval before commencing with IHC staining protocol.

Species	Human
Dilution info	1/1000
Notes	**For unpurified, use 1/100 - 1/250.** Perform heat-mediated antigen retrieval before commencing with IHC staining protocol.

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Rat
Dilution info	1/1000 - 1/10000
Notes	-

Species	Human
Dilution info	1/1000 - 1/10000
Notes	-

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1 µg/mL
Notes	This product gave a positive signal in THP-1 (-ve: HCT116) fixed with 4% formaldehyde (10 min).

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	-
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

ChIC/CUT&RUN-seq

Tested

Species	Human
Dilution info	5 µg
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

Target data

See full target information ERG

Function	Transcriptional regulator. May participate in transcriptional regulation through the recruitment of SETDB1 histone methyltransferase and subsequent modification of local chromatin structure.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot

Storage information	Avoid freeze / thaw cycle

Notes

Anti-ERG antibody [EPR3864] (ab92513) was developed by Abcam using patented rabbit monoclonal antibody technology and is validated for use in Flow Cyt (Intra), ICC/IF, IHC-P and WB.

Anti-ERG antibody [EPR3864] (ab92513) was first used in a scientific publication in 2010 and has been cited over 238 times in peer reviewed journals. It's performance in IHC in mouse and human samples is trusted by the scientific community.

Abcam's high quality manufacturing and validation processes ensure Anti-ERG antibody [EPR3864] (ab92513) has high sensitivity and specificity alongside high lot-to-lot consistency and reproducibility.

Anti-ERG antibody [EPR3864] (ab92513) has 17 independent reviews from customers.

Anti-ERG antibody [EPR3864] (ab92513) specifically detects ERG (UniProt ID: P11308; Molecular weight: 54kDa) and is sold in 100 µL and 1 mL selling sizes.

Conjugation-ready, carrier free format available for antibody clone EPR3864 - ab214796.

Antibody clone EPR3864 is also available pre-conjugated to a variety of labels for your convenience - HRP, Alexa Fluor® 647, Alexa Fluor® 488, PE, APC (ab195514, ab196149, ab196374, ab314261, ab315146).

ERG is a transcription factor involved in various cancers, including prostate cancer, where it is often overexpressed due to gene fusions. It plays a crucial role in tumor progression and metastasis, making it a potential target for therapeutic interventions.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	ERG or ETS-related gene is a transcription factor that belongs to the ETS (E-26 transformation specific) family. The ERG protein plays an important role in gene

	regulation involved in cell growth and differentiation. ERG has a molecular mass of approximately 54 kDa. It is expressed in a variety of tissues including the prostate endothelial cells and hematopoietic cells. Researchers often study ERG protein localization through techniques such as ERG IHC to better understand its expression across different tissues for example in the ERG color sections that highlight specific cellular contexts.
Biological function summary	ERG regulates processes critical for normal cellular functions such as proliferation apoptosis and differentiation. It functions as part of a larger transcriptional regulatory complex and interacts with other proteins to exert its effects on target genes. The ERG protein can affect angiogenesis and limb development through its regulation of vascular endothelial cells. By controlling such important cellular processes ERG influences both normal physiological functions and abnormal conditions when dysregulated.
Pathways	ERG is highly significant in the MAPK/ERK signaling pathway which is involved in cell division and survival. ERG interacts with other signaling proteins such as FLI1 and RUNX1 affecting cellular responses and contributing to the regulation of gene expression essential for development and homeostasis. Within the hematopoietic context ERG participates in the regulation of pathways that maintain stem cell populations making it important for normal blood cell development.
Associated diseases and disorders	ERG is prominently implicated in prostate cancer where gene fusions involving ERG lead to its overexpression and contribute to malignancy. The abnormal expression of ERG in endothelial cells can also have a role in vascular disorders. ERG's interplay with proteins like TMPRSS2 often seen in gene fusions in prostate cancer underlines its role in oncogenesis. Understanding how ERG regulates these pathways and its interactions with other proteins helps in developing targeted therapies for disorders involving ERG dysregulation.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


19 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ERG antibody [EPR3864] (ab92513)

Immunohistochemical analysis of paraffin embedded Human Prostatic adenocarcinoma stage 3 tissue using unpurified ab92513 showing +ve staining.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-ERG antibody [EPR3864] (ab92513)


ERG Flow Cytometry (Intracellular) staining using rabbit Anti-ERG antibody

Flow cytometry overlay histogram showing left THP-1 positive cells and right negative HCT116 stained with ab92513 (red line). The cells were fixed with 4% formaldehyde (10 min) and then permeabilised with 0.1% PBS-Triton X-100 for 15 min. The cells were then incubated in 1x PBS containing 10µg/ml human IgG and 10% normal goat serum to block FC receptors and non-specific protein-protein interaction followed by the antibody (ab92513) (1x 106 in 100µl at 0.04µg/ml (1/54000)) for 30min at 22°C.

The secondary antibody Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed was incubated at 1/4000 for 30min at 22°C

Isotype control antibody (black line) was Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control.

Acquisition of >5000 events were collected using a 50 mW Blue laser (488nm) and 525/40 bandpass filter.




Immunocytochemistry/ Immunofluorescence - Anti-ERG antibody [EPR3864] (ab92513)

ERG Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-ERG antibody

Immunofluorescence staining of THP-1 (human monocytic leukemia cell line) cells with purified ab92513 at a working dilution of 1/100, counter-stained with DAPI. The secondary antibody was Alexa Fluor® 488 goat anti-rabbit (ab150077), used at a dilution of 1/1000. ab7291, a mouse anti-tubulin antibody (1/1000), was used to stain tubulin along with ab150120 (Alexa Fluor® 594 goat anti-mouse, 1/1000), shown in the top right hand panel. The cells were fixed in 4% PFA and permeabilized using 0.1% Triton X 100. The negative controls are shown in bottom middle and right hand panels - for negative control 1, purified ab92513 was used at a dilution of 1/500


followed by an Alexa Fluor® 594 goat anti-mouse antibody (ab150120) at a dilution of 1/500. For negative control 2, ab7291 (mouse anti-tubulin) was used at a dilution of 1/500 followed by an Alexa Fluor® 488 goat anti-rabbit antibody (ab150077) at a dilution of 1/400.
Alexa Fluor® 488 (ab196374) and Alexa Fluor® 647 (ab196149) conjugated versions are available for this clone.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ERG antibody [EPR3864] (ab92513)

ERG Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-ERG antibody


Immunohistochemical staining of paraffin embedded human kidney with purified ab92513 at a working dilution of 1/1000. The secondary antibody used is HRP goat anti-rabbit IgG H&L (ab97051) at 1/500. The sample is counter-stained with hematoxylin. Antigen retrieval was perfomed using Tris-EDTA buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control, and is shown in the inset.



Flow Cytometry (Intracellular) - Anti-ERG antibody [EPR3864] (ab92513)

ERG Flow Cytometry (Intracellular) staining using rabbit Anti-ERG antibody

Intracellular Flow Cytometry analysis of THP-1 (human monocytic leukemia cell line) cells labeling ERG with purified ab92513 at 1/1000 dilution (1ug/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. A Goat anti rabbit IgG (Alexa Fluorr® 488) (ab150077) (1/2000 dilution) was used as the secondary antibody. Rabbit monoclonal IgG (Black) (ab172730) was used as the isotype control, Cell without incubation with primary antibody and secondary antibody (Blue) were used as the unlabeled control.



Immunocytochemistry/ Immunofluorescence - Anti-ERG antibody [EPR3864] (ab92513)

ERG Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-ERG antibody

ab92513 staining ERG in THP-1 cells, with negative expression in HCT116 cells. The cells were fixed with 4% formaldehyde (10 min), permeabilised with 0.1% Triton x-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab92513 at 1 µg/ml and ab7291, Mouse monoclonal [DM1A] to alpha Tubulin at 0.5 µg/ml. Cells were then incubated with ab150081, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 488), pre-adsorbed at 1/1000 dilution (shown in green) and ab150119, Goat polyclonal Secondary Antibody to Mouse IgG - H&L (Alexa Fluor® 647), pre-adsorbed at 1/1000 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue). Image was acquired with a confocal microscope (Leica-Microsystems TCS SP8) and a single confocal section is shown.





Image from Litovkin K et al. PLoS One. 2015;10(6):e0130651. Fig 5.; doi: 10.1371/journal.pone.0130651.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ERG antibody [EPR3864] (ab92513)

ERG and GSTP1 immunostainings of human prostate cancer samples using ab92513. Representative immunohistochemical images of prostate cancer samples are shown that were positive for ERG and negative for GSTP1 (A), positive for both ERG and GSTP1 (B), negative for both ERG and GSTP1(C), and negative for ERG and positive for GSTP1 (D). The internal staining control for ERG is the endothelium (arrows) and for GSTP1 the stromal and/or basal cells of normal prostate glands. N, normal prostate gland; S, Stroma; T, tumor gland. Scale bars equal 100µm



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ERG antibody [EPR3864] (ab92513)

Immunohistochemical analysis of formalin-fixed paraffin-embedded human kidney labelling ERG with ab92513 at a concentration of 1µg/ml. The immunostaining was performed on a Ventana DISCOVERY ULTRA (Roche Tissue Diagnostics) instrument with an OptiView DAB IHC Detection Kit. Heat mediated antigen retrieval was conducted for 32min with ULTRA cell conditioning solution (CC1 pH8.5). ab92513 anti ERG antibody was incubated at 37°C for 16min. Sections were counterstained with Hematoxylin II. Image inset shows absence of staining in secondary antibody only control.





Image from Kirby BJ et al. PLoS One. 2012;8(12):e83903. Fig 4.; doi: 10.1371/journal.pone.0035976.

Immunocytochemistry/ Immunofluorescence - Anti-ERG antibody [EPR3864] (ab92513)


Functional characterization and detection of genetic alterations in GEDI-captured cells. The TMPRSS2:ERG fusion protein is detected in GEDI-captured circulating tumor cells (CTCs) from a castrate-resistant prostate cancer (CRPC) patient. PSMA-captured CTCs were stained on the device with ab92513. Representative examples of PSMA+/CD45− CTCs are shown, two of which are positive for ERG. Scale bars: 10 microns.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ERG antibody [EPR3864] (ab92513)

ERG Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-ERG antibody

Formalin-fixed, paraffin-embedded mouse brain tissue stained for ERG using ab92513 at 1/200 dilution in immunohistochemical analysis. A horse radish peroxidase antibody was used as the secondary antibody.
Antigen Retrieval: 40x; Proteinase K antigen retrieval - 15 min at 37 C



Western blot - Anti-ERG antibody [EPR3864] (ab92513)

Blocking buffer: 5% NFDM/TBST
Dilution buffer: 5% NFDM/TBST
All lanes:
Western blot - Anti-ERG antibody [EPR3864] (ab92513) at 1/2000 dilution

Lane 1:
Jurkat (human T cell leukemia cell line from peripheral blood) cell lysate at 20 µg


Lane 2:
HeLa (human epithelial cell line from cervix adenocarcinoma) cell lysate at 20 µg

Secondary

All lanes:
HRP goat anti-rabbit IgG (H+L) at 1/1000 dilution


Predicted band size: 54 kDa

Observed band size: 55 kDa




ChIC/CUT&RUN sequencing - Anti-ERG antibody [EPR3864] (ab92513)

ChIC/CUT&RUN was performed using a pAG-MNase at a final concentration of 700 ng/mL, 2.5 x 10^5 HUVEC (human umbilical vein endothelial cell) cells and 5 µg of ab92513 [EPR3864]. The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control ab172730 is also shown.
The University of Geneva owns patents relevant to ChIC (Chromatin Immuno-Cleavage) methods.



ChIC/CUT&RUN sequencing - Anti-ERG antibody [EPR3864] (ab92513)


ChIC/CUT&RUN was performed using a pAG-MNase at a final concentration of 700 ng/mL, 2.5 x 10^5 HUVEC (human umbilical vein endothelial cell) cells and 5 µg of ab92513 [EPR3864]. The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control ab172730 is also shown.
The University of Geneva owns patents relevant to ChIC (Chromatin Immuno-Cleavage) methods.



ChIC/CUT&RUN sequencing - Anti-ERG antibody [EPR3864] (ab92513)

ChIC/CUT&RUN was performed using a pAG-MNase at a final concentration of 700 ng/mL, 2.5 x 10^5 HUVEC (human umbilical vein endothelial cell) cells and 5 µg of ab92513 [EPR3864]. The

resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control ab172730 is also shown.
The University of Geneva owns patents relevant to ChIC (Chromatin Immuno-Cleavage) methods.



Western blot - Anti-ERG antibody [EPR3864] (ab92513)

Blocking buffer: 5% NFDM/TBST
Dilution buffer: 5% NFDM/TBST
All lanes:
purified

Lane 1:
rat brain lysate at 20 µg

Lane 2:
rat heart lysate at 20 µg


Lane 3:
RAW 264.7 (mouse macrophage cell line transformed with Abelson murine leukemia virus) cell lysate at 20 µg

Secondary

All lanes:
HRP goat anti-rabbit IgG (H+L) at 1/1000 dilution


Predicted band size: 54 kDa

Observed band size: 55 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ERG antibody [EPR3864] (ab92513)

Tissue Microarrays stained for Anti-ERG antibody [EPR3864] using ab92513 in immunohistochemical analysis. This table provides a detailed overview of positive (tick mark) and negaive (cross mark) staining per sample type tested. The section was incubated with ab92513 for 30 mins at room temperature followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Western blot - Anti-ERG antibody [EPR3864] (ab92513)


All lanes:
Western blot - Anti-ERG antibody [EPR3864] (ab92513) at 1/1000 dilution

All lanes:
Jurkat (human T cell leukemia cell line from peripheral blood) cell lysate at 10 µg

Secondary

All lanes:
HRP labelled Goat anti-Rabbit at 1/2000 dilution

Predicted band size: 54 kDa



Western blot - Anti-ERG antibody [EPR3864] (ab92513)

Lanes 1 and 3:
Western blot - Anti-ERG antibody [EPR3864] (ab92513) at 1/250 dilution

Lanes 2 and 4:
Western blot - Anti-ERG antibody [EPR3864] (ab92513) at 1/1000 dilution

Lane 1:
Jurkat (human T cell leukemia cell line from peripheral blood) Whole Cell Lysate at 10 µg

Lanes 2 - 4:
Jurkat Whole Cell Lysate at 10 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/10000 dilution


Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 54 kDa

Observed band size: 55 kDa

Exposure time: 12min



OI-RD Scanning - Anti-ERG antibody [EPR3864] (ab92513)

We have systematically measured KD (the equilibrium dissociation constant between the antibody and its antigen), of more than 840 recombinant antibodies to assess not only their individual KD values but also to see the average affinity of antibody. Based on the comparison with published literature values for mouse monoclonal antibodies, Recombinant antibodies appear to be on average 1-2 order of magnitude higher affinity.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.