abcam

ab92536

Anti-MMP2 antibody [EPR1184]

Anti-MMP2 antibody [EPR1184] (ab92536) is a rabbit monoclonal antibody detecting MMP2 in **Western Blot, Flow Cytometry (Intra), ICC/IF**. Suitable for **Human, Mouse, Rat**.

- Biophysical QC for unrivalled batch-batch consistency
- Over 370 publications
- Trusted since 2010

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR1184
Purification technique	Affinity purification Protein A
Specificity	In Western Blot, this product typically gives a weaker signal in some hepatocarcinoma like HepG2, MHCC97L (PMID: 33184263) et al. Please use recommended positive control when testing these cells. Compared with ab92536, ab181286 has higher sensitivity. We recommend ab181286 as an alternative for testing MMP2 in western blot. This antibody shows low affinity in detecting mouse liver and HepG2 lysates which are positive for MMP2 reported by PMID 24096707 and 24297510.
Concentration	1.68 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Mouse
Dilution info	1/1000 - 1/5000
Notes	For Lysate preparation protocol, please refer to the protocol here. Compared with ab92536, ab181286 has higher sensitivity. We recommend ab181286 as an alternative for testing MMP2 in western blot.

Species	Rat
Dilution info	1/1000 - 1/5000
Notes	For Lysate preparation protocol, please refer to the protocol here. Compared with ab92536, ab181286 has higher sensitivity. We recommend ab181286 as an alternative for testing MMP2 in western blot.

Species	Human
Dilution info	1/1000 - 1/5000
Notes	For Lysate preparation protocol, please refer to the protocol here. Compared with ab92536, ab181286 has higher sensitivity. We recommend ab181286 as an alternative for testing MMP2 in western blot.

IHC-P

Not recommended

Species	Mouse, Rat, Human
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human

Dilution info	1/250
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/400
Notes	For unpurified, use 1/70. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information MMP2

Function Ubiquitinous metalloproteinase that is involved in diverse functions such as remodeling of the vasculature, angiogenesis, tissue repair, tumor invasion, inflammation, and atherosclerotic plaque rupture. As well as degrading extracellular matrix proteins, can also act on several nonmatrix proteins such as big endothelial 1 and beta-type CGRP promoting vasoconstriction. Also cleaves KISS at a Gly-|-Leu bond. Appears to have a role in myocardial cell death pathways. Contributes to myocardial oxidative stress by regulating the activity of GSK3beta. Cleaves GSK3beta in vitro. Involved in the formation of the fibrovascular tissues in association with MMP14. PEX, the C-terminal non-catalytic fragment of MMP2, possesses anti-angiogenic and anti-tumor properties and inhibits cell migration and cell adhesion to FGF2 and vitronectin. Ligand for integrinv/beta3 on the surface of blood vessels. Isoform 2. Mediates the proteolysis of CHUK/IKKA and initiates a primary innate immune response by inducing mitochondrial-nuclear

stress signaling with activation of the pro-inflammatory NF-kappaB, NFAT and IRF transcriptional pathways.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Stable for 12 months at -20°C

Notes

Product Specifications
Anti-MMP2 antibody [EPR1184] (ab92536) was developed by Abcam using patented rabbit monoclonal antibody technology and is validated for use in Flow Cyt (Intra), ICC/IF, WB in human, mouse, rat samples.
Anti-MMP2 antibody [EPR1184] (ab92536) specifically detects MMP2 (UniProt ID: P08253; Molecular weight: 62kDa) and is sold in 100 µL and 1 mL selling sizes.

Quality and Validation
Abcam's high quality manufacturing and validation processes ensure Anti-MMP2 antibody [EPR1184] (ab92536) has high sensitivity and specificity alongside high lot-to-lot consistency and reproducibility.
Anti-MMP2 antibody [EPR1184] (ab92536) has been cited over 376 times in peer reviewed journals and is trusted by the scientific community.
Anti-MMP2 antibody [EPR1184] (ab92536) has 7 independent reviews from customers.

Related Products
Antibody clone EPR1184 is also available pre-conjugated to a variety of labels for your convenience - Alexa Fluor® 488, Alexa Fluor® 647, APC, PE, Alexa Fluor® 594, Alexa Fluor® 568, Alexa Fluor® 555 (ab237473, ab237474, ab310884, ab310940, ab311718, ab312994, ab313199).

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity

- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The MMP-2 protein also known as matrix metalloproteinase-2 or gelatinase A is an enzyme involved in the breakdown of extracellular matrix components. It plays a critical role in tissue remodeling and cell migration. Comprised of a molecular weight of approximately 72 kDa this metalloproteinase is secreted as an inactive proenzyme that requires activation. MMP2 is expressed in various tissues including the brain heart and blood vessels where it contributes to normal physiological processes and pathological conditions.
Biological function summary	Matrix metalloproteinase-2 is mainly involved in the degradation of type IV and V collagens gelatin and fibronectin. As part of the metalloproteinase family it works alongside other MMPs to maintain tissue homeostasis and repair. MMP-2 forms part of a complex network that ensures the timely degradation of matrix components balancing synthesis and breakdown. It remains regulated by tissue inhibitors of metalloproteinases (TIMPs) preventing excessive degradation that could lead to tissue damage.
Pathways	MMP-2 plays a significant role within the extracellular matrix (ECM) remodeling and angiogenesis pathways. It interacts with various proteins including integrins and TIMP-2 to modulate cellular behaviors such as migration and invasion. MMP-2 contributes to processes like wound healing and embryonic development through its involvement in ECM degradation and new tissue formation.
Associated diseases and disorders	Matrix metalloproteinase-2 is linked to cancer progression and cardiovascular diseases. In cancer abnormal MMP-2 activity facilitates tumor invasion and metastasis by breaking down matrix barriers. Increased MMP-2 expression associates with poor prognosis in cancers like breast and prostate. In cardiovascular diseases such as atherosclerosis it contributes to plaque destabilization and vascular remodeling. The imbalance in MMP-2 activity and its regulation by proteins like TIMP-1 are involved in the pathology of these disorders.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

9 product images



Western blot - Anti-MMP2 antibody [EPR1184] (ab92536)

Blocking and diluting buffer and concentration: 5% NFDM /TBST. ab181602 was used as GAPDH loading control. Compared with ab92536, ab181286 has higher sensitivity. We recommend ab181286 as an alternative for testing MMP2 in western blot.
Lanes 1 - 5:
Western blot - Anti-MMP2 antibody [EPR1184] (ab92536) at 1/1000 dilution

Lanes 1 - 5:
Western blot - Anti-MMP2 antibody [EPR17003-25] (ab181286)

Lane 1:
HT-1080 (Human fibrosarcoma epithelial cell) whole cell lysate at 20 µg

Lane 2:
Untreated HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysate at 20 µg

Lane 3:
HepG2 (Human hepatocellular carcinoma epithelial cell) treated with 300ng/ml BFA for 24 hours whole cell lysate at 20 µg

Lane 4:
Huh7 (Human hepatocellular carcinoma epithelial cell) whole cell lysate at 20 µg


Lane 5:
U-87 MG (Human glioblastoma-astrocytoma epithelial cell) whole cell lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

Predicted band size: 73 kDa


Exposure time: 60s



Flow Cytometry (Intracellular) - Anti-MMP2 antibody [EPR1184] (ab92536)

MMP2 Flow Cytometry (Intracellular) staining using rabbit Anti-MMP2 antibody


Intracellular Flow Cytometry analysis of PC-3 (human prostate adenocarcinoma) cells labeling MMP2 with purified ab92536 at 1/180 dilution (10ug/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. A Goat anti rabbit IgG (Alexa Fluorr® 488) (1/2000 dilution) was used as the secondary antibody. Rabbit monoclonal IgG (Black) was used as the isotype control, cells without incubation with primary antibody and secondary antibody (Blue) were used as the unlabeled control.



Immunocytochemistry/ Immunofluorescence - Anti-MMP2 antibody [EPR1184] (ab92536)

MMP2 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-MMP2 antibody


Immunofluorescence staining of PC-3 cells with purified ab92536 at a working dilution of 1/250 counter-stained with DAPI. The secondary antibody was Alexa Fluor® 488 goat anti-rabbit (ab150077) used at a dilution of 1/1000. ab7291 a mouse anti-tubulin antibody (1/1000) was used to stain tubulin along with ab150120 (Alexa Fluor® 594 goat anti-mouse 1/1000) shown in the top right hand panel. The cells were fixed in 4% PFA and permeabilized using 0.1% Triton X 100. The negative controls are shown in bottom middle and right hand panels - for negative control 1 purified ab92536 was used at a dilution of 1/500 followed by an Alexa Fluor® 594 goat anti-mouse antibody (ab150120) at a dilution of 1/500. For negative control 2 ab7291 (mouse anti-tubulin) was used at a dilution of 1/500 followed by an Alexa Fluor® 488 goat anti-rabbit antibody (ab150077) at a dilution of 1/400.



Flow Cytometry (Intracellular) - Anti-MMP2 antibody [EPR1184] (ab92536)

MMP2 Flow Cytometry (Intracellular) staining using rabbit Anti-MMP2 antibody

Overlay histogram showing HeLa cells fixed in 4% PFA and stained with purified ab92536 at a dilution of 1 in 400 (red line). The secondary antibody used was FITC goat anti-rabbit at a dilution of 1 in 500. Rabbit monoclonal IgG was used as an isotype control (black line) and cells incubated in the absence of both primary and secondary antibody were used as a negative control (blue line).



Western blot - Anti-MMP2 antibody [EPR1184] (ab92536)

MMP2 Western blot staining using rabbit Anti-MMP2 antibody

The 72 KDa band is pro-MMP2 and the 69 KDa band is active-MMP2 reported by PMID 11489818 and 22190701.
Blocking and diluting buffer and concentration: 5% NFDM/TBST.

This antibody shows low affinity in detecting mouse liver and HepG2 lysates which are positive for MMP2 reported by PMID 24096707 and 24297510.

All lanes:
Western blot - Anti-MMP2 antibody [EPR1184] (ab92536) at 1/1000 dilution

Lane 1:

L6 (Rat skeletal muscle myoblast) whole cell lysates prepared in 1%SDS Hot lysis method at 20 µg

Lane 2:
Mouse liver lysates prepared in RIPA lysis method at 20 µg

Lane 3:
Mouse liver lysates prepared in 1%SDS Hot lysis method at 20 µg

Lane 4:
Raw264.7 (Mouse Abelson murine leukemia virus-induced tumor macrophage) whole cell lysates prepared in RIPA lysis method at 20 µg

Lane 5:
Raw264.7 (Mouse Abelson murine leukemia virus-induced tumor macrophage) whole cell lysates prepared in 1%SDS Hot lysis method at 20 µg

Lane 6:
HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysates prepared in RIPA lysis method at 20 µg

Lane 7:
HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysates prepared in 1%SDS Hot lysis method at 20 µg


Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 73 kDa

Observed band size: 69 kDa, 72 kDa

Exposure time: 10s



Western blot - Anti-MMP2 antibody [EPR1184] (ab92536)

Blocking and diluting buffer and concentration: 5% NFDM/TBST. ab181602 was used as a GAPDH loading control. Compared with ab92536, ab181286 has higher sensitivity. We recommend ab181286 as an alternative for testing MMP2 in western blot.
Lanes 1 - 3:
Western blot - Anti-MMP2 antibody [EPR1184] (ab92536) at 1/1000 dilution

Lanes 1 - 3:
Western blot - Anti-MMP2 antibody [EPR17003-25] (ab181286)

Lane 1:
Human plasma tissue lysate at 20 µg

Lane 2:
Human brain tissue lysate at 20 µg

Lane 3:
Human breast tissue lysate at 20 µg


Secondary

All lanes:
Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

Predicted band size: 73 kDa

Observed band size: 69 kDa, 72 kDa

Exposure time: 60s



Western blot - Anti-MMP2 antibody [EPR1184] (ab92536)

MMP2 Western blot staining using rabbit Anti-MMP2 antibody

All lanes:
Western blot - Anti-MMP2 antibody [EPR1184] (ab92536) at 1/1000 dilution

Lane 1:
L6 cell lysate at 10 µg


Lane 2:
Fetal heart lysate at 10 µg

Lane 3:
NIH/3T3 cell lysate at 10 µg

Secondary

All lanes:
HRP labelled goat anti-rabbit at 1/2000 dilution

Predicted band size: 73 kDa



Western blot - Anti-MMP2 antibody [EPR1184] (ab92536)

MMP2 Western blot staining of human skin using rabbit Anti-MMP2 antibody

Blocking buffer: 5% NFDM/TBST
Dilution buffer: 5% NFDM/TBST
72kDa: propeptide; 64kDa: active form

All lanes:
Western blot - Anti-MMP2 antibody [EPR1184] (ab92536) at 1/5000 dilution


All lanes:
human skin at 10 µg

Secondary

All lanes:
HRP goat anti-rabbit IgG (H+L) at 1/1000 dilution

Predicted band size: 73 kDa

Observed band size: 64 kDa, 72 kDa



Western blot - Anti-MMP2 antibody [EPR1184] (ab92536)

MMP2 Western blot staining using rabbit Anti-MMP2 antibody

Blocking buffer: 5% NFDM/TBST
Dilution buffer: 5% NFDM/TBST
72kDa: propeptide; 64kDa: active form

All lanes:
Western blot - Anti-MMP2 antibody [EPR1184] (ab92536) at 1/10000 dilution

Lane 1:
L6 cell lysate at 20 µg

Lane 2:
NIH/3T3 cell lysate at 20 µg

Secondary

All lanes:
HRP goat anti-rabbit IgG (H+L) at 1/1000 dilution

Predicted band size: 73 kDa

Observed band size: 64 kDa, 72 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.