abcam

ab95540

PE/Cy5® Anti-CD62L antibody [DREG-56]

Anti-CD62L antibody [DREG-56] conjugated to PE/Cy5® - suitable for flow cytometry, and reacts with human samples. Clone [DREG-56] is a mouse monoclonal antibody and one of the most commonly used clones for flow cytometry.

Design your panel with our interactive flow cytometry panel builder. Click here.

Key facts

Isotype	IgG1
Host species	Mouse
Conjugation	PE/Cy5®
Excitation/Emission	Ex: 496nm, Em: 667nm
Storage buffer	pH: 7.2 Preservative: 0.09% Sodium azide Constituents: PBS, 0.87% Sodium chloride, 0.2% BSA
Form	Liquid
Clonality	Monoclonal
Clone number	DREG-56
Purification technique	Affinity purification Protein G
Light chain type	kappa

Reactivity data

Flow Cyt

Tested

Species	Human
Dilution info	5 µL for 10^6 Cells

Notes	ab67435 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

Target data

See full target information SELL

Function	Calcium-dependent lectin that mediates cell adhesion by binding to glycoproteins on neighboring cells (PubMed:12403782, PubMed:28011641, PubMed:28489325). Mediates the adherence of lymphocytes to endothelial cells of high endothelial venules in peripheral lymph nodes. Promotes initial tethering and rolling of leukocytes in endothelia (PubMed:12403782, PubMed:28011641).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	CD62L also known as L-selectin is a cell adhesion molecule with a mass of approximately 74 kDa. It is part of the L-selectin family and is found on the surface of leukocytes including T cells B cells neutrophils and monocytes. Mouse models frequently study CD62L under the name "MEL-14". CD62L plays an important role in mediating the adhesion and rolling of leukocytes on blood vessel endothelial cells during the immune response. Its structure allows it to interact specifically with carbohydrates on endothelial cells facilitating the initial steps of leukocyte extravasation into tissues.
Biological function summary	CD62L is involved in the migration of leukocytes into lymphoid tissues and sites of inflammation. It is not known to be part of a larger protein complex; rather it functions independently on the cell membrane. CD62L also assists in the homing of naive lymphocytes to peripheral lymph nodes. The molecule regulates how immune cells enter tissues through high endothelial venules acting as a critical entry point for leukocyte trafficking.
Pathways	CD62L contributes significantly to the immune trafficking pathway and leukocyte adhesion cascade. It works closely with other adhesion molecules like CD44 and CD54 (ICAM-1) which are important within the context of cellular responses during inflammation. CD62L's function in these pathways ensures that immune cells are

	properly directed to areas where they are needed most whether that is secondary lymphoid organs or inflamed tissues.
Associated diseases and disorders	CD62L expression and function have been associated with inflammatory conditions and certain immune-related pathologies such as multiple sclerosis and rheumatoid arthritis. Altered CD62L expression can affect lymphocyte homing potentially playing a role in the pathogenesis of these conditions. The role of CD62L is also linked to proteins like CD44 which can modulate cell interactions and have overlapping functions in disease mechanisms influencing disease progression or resolution.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

1 product image



Flow Cytometry - PE/Cy5® Anti-CD62L antibody [DREG-56] (ab95540)

CD62L Flow Cytometry staining using mouse Anti-CD62L antibody

Staining of normal Human peripheral blood cells with PE-Cy5 Mouse IgG1 kappa Isotype Control (open histogram) or ab95540 (filled histogram).
Cells in the lymphocyte gate were used for analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.