abcam

ab95816

Biotin Anti-Integrin beta 7 antibody [FIB504]

Anti-Integrin beta 7 antibody [FIB504] conjugated to Biotin - suitable for flow cytometry, ICC, and reacts with mouse, human samples. Clone [FIB504] is a rat monoclonal antibody cited in 45 publications.

Key facts

Isotype	IgG2a
Host species	Rat
Conjugation	Biotin
Storage buffer	pH: 7.2 Preservative: 0.09% Sodium azide
Form	Liquid
Clonality	Monoclonal
Immunogen	Cell preparation containing Itgb7 protein. The exact immunogen used to generate this antibody is proprietary information. Database link P26011
Clone number	FIB504
Purification technique	Affinity purification Protein G
Light chain type	kappa
Concentration	0.5 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC

Tested

Species	Mouse

Dilution info	-
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt

Tested

Species	Human
Dilution info	0.125 µg for 10^5 Cells
Notes	Range from 10^5 to 10^8 cells/test in a final volume of 100 µl. ab18445 - Rat monoclonal IgG2a, is suitable for use as an isotype control with this antibody.

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information Itgb7

Function	Integrin ITGA4/ITGB7 (alpha-4/beta-7) (Peyer patches-specific homing receptor LPAM-1) is an adhesion molecule that mediates lymphocyte migration and homing to gut-associated lymphoid tissue (GALT). Integrin ITGA4/ITGB7 interacts with the cell surface adhesion molecules MADCAM1 which is normally expressed by the vascular endothelium of the gastrointestinal tract. Also interacts with VCAM1 and fibronectin, an extracellular matrix component. It recognizes one or more domains within the alternatively spliced CS-1 region of fibronectin. Interactions involve the tripeptide L-D-T in MADCAM1, and L-D-V in fibronectin. Integrin ITGAE/ITGB7 (alpha-E/beta-7, HML-1) is a receptor for E-cadherin.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Storage information	Avoid freeze / thaw cycle

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Integrin beta 7 also known as beta-7 integrin beta 7 or β7 is a protein encoded by the ITGB7 gene. It forms a heterodimer most commonly with integrin alpha 4 (forming α4β7 integrin) or integrin alpha E (forming αEβ7 integrin). The molecular mass of integrin beta 7 is approximately 84 kDa. It is expressed in the intestinal mucosa particularly associated with immune cells such as T lymphocytes and dendritic cells. Integrin beta 7 plays an important role in cell adhesion and migration by binding to its ligands which are present on endothelial cells in the gut.
Biological function summary	Integrin beta 7 facilitates cell adhesion and signaling processes important for immune surveillance and inflammation regulation. As part of heterodimer complexes like α4β7 and αEβ7 integrins it mediates adhesion of lymphocytes to mucosal tissues. This interaction is necessary for directing lymphocytes to sites of inflammation in the intestinal tract. The α4β7 integrin particularly binds to the addressin MAdCAM-1 which is expressed on high endothelial venules in the gut-associated lymphoid tissues.
Pathways	Integrin beta 7 is involved in both integrin-mediated signaling and leukocyte extravasation pathways. These pathways regulate the localization and migration of immune cells to target tissues which is essential for efficient immune responses. The protein works together with integrin alpha 4 and alpha E components to facilitate these processes. Integrin signaling activates various intracellular cascades including those involving kinases that modulate cell adhesion and migration.
Associated diseases and disorders	Integrin beta 7 has significant implications in inflammatory bowel disease (IBD) and other immune-mediated conditions like celiac disease. Elevated expression of integrin beta 7 particularly in the α4β7 complex associates with the overactive lymphocyte trafficking found in IBD. As a result therapies targeting this integrin like vedolizumab aim to modulate lymphocyte migration. The involvement of proteins such as MAdCAM-1 in this integrin's function links it to the pathophysiology of these diseases.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


3 product images



Flow Cytometry - Biotin Anti-Integrin beta 7 antibody [FIB504] (ab95816)

Integrin beta 7 Flow Cytometry staining using rat Anti-Integrin beta 7 antibody


Staining of normal Human peripheral blood cells with Anti-Human CD3 APC and 0.125 µg of Rat IgG2a kappa Isotype Control Biotin. Cells in the lymphocyte gate were used for analysis.



Flow Cytometry - Biotin Anti-Integrin beta 7 antibody [FIB504] (ab95816)

Integrin beta 7 Flow Cytometry staining using rat Anti-Integrin beta 7 antibody

Staining of normal Human peripheral blood cells with 0.06 µg of ab95816 followed by Streptavidin PE. Cells in the lymphocyte gate were used for analysis.



This image is courtesy of an anonymous customer review.

Immunocytochemistry - Biotin Anti-Integrin beta 7 antibody [FIB504] (ab95816)

Integrin beta 7 Immunocytochemistry staining using rat Anti-Integrin beta 7 antibody

ICC/IF image of Integrin beta 7 staining on RAW264.7 cell line using ab95816 (1:300). The cells were fixed using formaldehyde and permeabilized using 0.2% Triton X-100. The cells were blocked using 10% goat serum for 1 hour at 20°C. The cells were incubated with ab95816 (1:300) for 1 hour at 20°C. The staining was detected using Streptavedin conjugated to Alexo Fluor 647 (1:350)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.