abcam

ab97613

Anti-EGFR (phospho Y845) antibody [EPR2149Y]

Lab Essentials | Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR2149Y
Purification technique	Affinity purification Protein A
Concentration	0.415 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Dot

Tested

Species	Synthetic peptide
Dilution info	-
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Human
Dilution info	1/500
Notes	-

Not recommended

Species	Synthetic peptide
Dilution info	-
Notes	-

IHC-P

Not recommended

Species	Human, Synthetic peptide
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/100 - 1/250
Notes	-

Not recommended

Species	Synthetic peptide
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/400
Notes	-

Not recommended

Species	Synthetic peptide
Dilution info	-
Notes	-

Target data

See full target information EGFR pY845

Function

Receptor tyrosine kinase binding ligands of the EGF family and activating several signaling cascades to convert extracellular cues into appropriate cellular responses (PubMed:10805725, PubMed:27153536, PubMed:2790960, PubMed:35538033). Known ligands include EGF, TGFA/TGF-alpha, AREG, epigen/EPGN, BTC/betacellulin, epiregulin/EREG and HBEGF/heparin-binding EGF (PubMed:12297049, PubMed:15611079, PubMed:17909029, PubMed:20837704, PubMed:27153536, PubMed:2790960, PubMed:7679104, PubMed:8144591, PubMed:9419975). Ligand binding triggers receptor homo- and/or heterodimerization and autophosphorylation on key cytoplasmic residues. The phosphorylated receptor recruits adapter proteins like GRB2 which in turn activates complex downstream signaling cascades. Activates at least 4 major downstream signaling cascades including the RAS-RAF-MEK-ERK, PI3 kinase-AKT, PLCgamma-PKC and STATs modules (PubMed:27153536). May also activate the NF-kappa-B signaling cascade (PubMed:11116146). Also directly phosphorylates other proteins like RGS16, activating its GTPase activity and probably coupling the EGF receptor signaling to the G protein-coupled receptor signaling (PubMed:11602604). Also phosphorylates MUC1 and increases its interaction with SRC and CTNNB1/beta-catenin (PubMed:11483589). Positively regulates cell migration via interaction with CCDC88A/GIV which retains EGFR at the cell membrane following ligand stimulation, promoting EGFR signaling which triggers cell migration (PubMed:20462955). Plays a role in enhancing learning and memory

performance (By similarity). Plays a role in mammalian pain signaling (long-lasting hypersensitivity) (By similarity). Isoform 2 may act as an antagonist of EGF action. (Microbial infection) Acts as a receptor for hepatitis C virus (HCV) in hepatocytes and facilitates its cell entry. Mediates HCV entry by promoting the formation of the CD81-CLDN1 receptor complexes that are essential for HCV entry and by enhancing membrane fusion of cells expressing HCV envelope glycoproteins.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Species reactivity
Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	EGFR or Epidermal Growth Factor Receptor is a transmembrane glycoprotein that acts as a receptor for members of the epidermal growth factor family. Known alternatively as ErbB1 or HER1 this receptor has an approximate molecular weight of 170 kDa. EGFR is expressed in various cell types notably on epithelial cells and can influence multiple cellular processes through its kinase activity. It participates in the regulation of cell growth multiplication and survival by activating its kinase domain upon ligand binding.
Biological function summary	The EGFR protein plays an important role in cellular communication and signaling processes. EGFR pairs with other receptor family members to form active dimers or even higher-order complexes which in turn initiate intracellular signaling cascades. Through these complexes EGFR influences many processes including cell differentiation and repair. This function of EGFR makes it an integral part of mammalian biology affecting how cells respond to their environment by mediating changes in gene expression.
Pathways	EGFR is a central player in the MAPK and PI3K/Akt signaling pathways. Alongside other protein partners like KRAS and PI3 kinase it contributes to transmitting signals from the cell surface to the nucleus affecting gene transcription and cell behavior. These pathways are important for normal cell growth and division and aberrations in these pathways can lead to excessive or insufficient cell proliferation.
Associated diseases and disorders	EGFR is pertinent to cancer biology including non-small cell lung cancer and glioblastoma where mutations or overexpression of the receptor frequently occur. It connects to proteins such as PTEN and BRAF which influence tumor progression and response to targeted therapies. EGFR's involvement in these disorders highlights its significance as a therapeutic target since it can be manipulated to alter disease progression.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

5 product images



Immunocytochemistry/ Immunofluorescence - Anti-EGFR (phospho Y845) antibody [EPR2149Y] (ab97613)


EGFR (phospho Y845) Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-EGFR (phospho Y845) antibody

Immunofluorescent analysis of 4% paraformaldehyde-fixed,0.1% Triton X-100 permeabilized A431 (Human epidermoid carcinoma cell line) cell lines labeling EGFR with ab97613 (purified) at 1/50 dilution,followed by Goat anti-Rabbit IgG H & L (Alexa Fluor® 488) (ab150077) secondary antibody at 1/1000 dilution
Cells were counterstained with ab195889 (Anti-alpha Tubulin antibody [DM1A]) - Microtubule Marker (Alexa Fluor® 594)

The nuclear counterstain is DAPI (blue).

The green staining on the membrane was increased in the EGF (100ng/ml, 10min) treated A431 cells when compared with A431 cells without treatment. After LP treatment, the green signaling was obviously decreased.


For the pan antibody, there was no great difference after EGF (100ng/ml, 10min) or EGF (100ng/ml, 10min) + LP treatment. The data showed mostly membranous staining.



Flow Cytometry (Intracellular) - Anti-EGFR (phospho Y845) antibody [EPR2149Y] (ab97613)

EGFR (phospho Y845) Flow Cytometry (Intracellular) staining using rabbit Anti-EGFR (phospho Y845) antibody

Intracellular Flow Cytometry analysis of A431 (Human epidermoid carcinoma epithelial cell) treated with 100ng/ml EGF for 10 minutes cells labeling EGFR with purified ab97613 at 1/400 dilution (1µg/ml) (red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluorr® 488, ab150077) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue). Untreated A431 cells (Green).



Western blot - Anti-EGFR (phospho Y845) antibody [EPR2149Y] (ab97613)

EGFR (phospho Y845) Western blot staining using rabbit Anti-EGFR (phospho Y845) antibody

Blocking and dilution buffer: 5% NFDM/TBST.
All lanes:
Western blot - Anti-EGFR (phospho Y845) antibody [EPR2149Y] (ab97613) at 1/1500 dilution

Lane 1:
Untreated A431 (Human epidermoid carcinoma cell line) whole cell lysates at 15 µg

Lane 2:
A431 (Human epidermoid carcinoma cell line) treated with 100 ng/ml EGF for 10 minutes whole cell lysates at 15 µg


Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 134 kDa

Observed band size: 175 kDa

Exposure time: 30s




Western blot - Anti-EGFR (phospho Y845) antibody [EPR2149Y] (ab97613)

All lanes:
Western blot - Anti-EGFR (phospho Y845) antibody [EPR2149Y] (ab97613) at 1/500 dilution

Lane 1:
A431 cell lysates, untreated at 10 µg

Lane 2:
A431 cell lysates, treated with EGF at 10 µg

Predicted band size: 134 kDa



Dot Blot - Anti-EGFR (phospho Y845) antibody [EPR2149Y] (ab97613)

Dot Blot analysis of Lane 1: EGFR (pY845) phospho peptide and Lane 2: EGFR non-phospho peptide labeling EGFR (phospho Y845) with ab97613 (unpurified) at 1/1000 dilution. 5% NFDM/TBST was used as the diluting and blocking buffer. ab97051 Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated was used as the secondary antibody at 1/100000 dilution. Exposure time: 3 minutes.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.