abcam

ab9809

Anti-TNF alpha antibody [NF-7]

Lab Essentials

Key facts

Isotype	IgG1
Host species	Mouse
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.1% Sodium azide
Form	Liquid
Clonality	Monoclonal
Immunogen	Recombinant Full Length Protein corresponding to Human TNF. Database link P01375
Clone number	NF-7
Purification technique	Affinity purification Protein A
Light chain type	unknown
Concentration	1 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Flow Cyt

Tested

Species	Human
Dilution info	2 µg for 10^6 Cells

Notes	ab170190 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

Target data

See full target information TNF

Function	Cytokine that binds to TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. It is mainly secreted by macrophages and can induce cell death of certain tumor cell lines. It is potent pyrogen causing fever by direct action or by stimulation of interleukin-1 secretion and is implicated in the induction of cachexia, Under certain conditions it can stimulate cell proliferation and induce cell differentiation. Impairs regulatory T-cells (Treg) function in individuals with rheumatoid arthritis via FOXP3 dephosphorylation. Up-regulates the expression of protein phosphatase 1 (PP1), which dephosphorylates the key 'Ser-418' residue of FOXP3, thereby inactivating FOXP3 and rendering Treg cells functionally defective (PubMed:23396208). Key mediator of cell death in the anticancer action of BCG-stimulated neutrophils in combination with DIABLO/SMAC mimetic in the RT4v6 bladder cancer cell line (PubMed:16829952, PubMed:22517918, PubMed:23396208). Induces insulin resistance in adipocytes via inhibition of insulin-induced IRS1 tyrosine phosphorylation and insulin-induced glucose uptake. Induces GKAP42 protein degradation in adipocytes which is partially responsible for TNF-induced insulin resistance (By similarity). Plays a role in angiogenesis by inducing VEGF production synergistically with IL1B and IL6 (PubMed:12794819). Promotes osteoclastogenesis and therefore mediates bone resorption (By similarity). The TNF intracellular domain (ICD) form induces IL12 production in dendritic cells.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

This antibody can be used in two-site immunoassay for the determination of human TNF alpha. It recognizes a different epitope than the monoclonal antibody ab9813. The two-site immunoassay can be used to determine TNF alpha concentration in the cultures of human peripheral blood mononuclear cells. Can also be used to neutralize TNF alpha cytotoxic activity in vitro. At a concentration of 2.5 µg/ml, antibody ab9809 causes 50% inhibition of TNF alpha mediated cytotoxicity on L929 cells.
This product was changed from ascites to tissue culture supernatant on 28/11/2017. Lot numbers higher than GR151594-8 will be from tissue culture supernatant. Please note that the dilutions may need to be adjusted accordingly.

This product was changed from ascites to tissue culture supernatant on 28/11/2017. Lot numbers higher than GR151594-8 will be from tissue culture supernatant. Please note that the dilutions may need to be adjusted accordingly.

This antibody can be used in two-site immunoassay for the determination of human TNF alpha. It recognizes a different epitope than the monoclonal antibody ab9813. The two-site immunoassay can be used to determine TNF alpha concentration in the cultures of human peripheral blood mononuclear cells. Can also be used to neutralize TNF alpha cytotoxic activity in vitro. At a concentration of 2.5 µg/ml, antibody ab9809 causes 50% inhibition of TNF alpha mediated cytotoxicity on L929 cells.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

1 product image



Flow Cytometry - Anti-TNF alpha antibody [NF-7] (ab9809)

TNF alpha Flow Cytometry staining using mouse Anti-TNF alpha antibody

Overlay histogram showing THP1 cells stained with ab9809 (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab9809, 2µg/1x10^6 cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) (ab96879) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG1 [ICIGG1] (ab91353, 2µg/1x10^6 cells) used under the same conditions. Acquisition of >5,000 events was performed.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.