

Product datasheet

Anti-GFAP antibody [GF5] ab10062

★★★★★ 22 Abreviews 131 References 7 Images

Overview

Product name	Anti-GFAP antibody [GF5]
Description	Mouse monoclonal [GF5] to GFAP
Host species	Mouse
Specificity	There is no cross-reactivity with other neurospecific proteins.
Tested applications	Suitable for: IHC-P, WB, ELISA, IHC-FoFr, IHC-Fr, ICC/IF, Flow Cyt, ICC
Species reactivity	Reacts with: Mouse, Rat, Human
Immunogen	This clone has been derived from hybridization of Sp2/0 myeloma cells with spleen cells of Balb/c mice immunised with purified glial fibrillary acidic protein from human brain.
Positive control	normal adult rat brain: lateral ventricle IHC-P: FFPE human hippocampus normal. IHC-P: FFPE rat brain normal.
General notes	<p>Abcam is committed to meeting high quality standards of ethical manufacturing and has decided to discontinue this product by June 2020 as it has been generated by the ascites method. We are sorry for any inconvenience this may cause. We suggest ab4674, ab68428 or ab7260 as possible replacements.</p> <p>Concentration varies from lot to lot and can be provided on request.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot. Store at +4°C. Do Not Freeze.
Storage buffer	pH: 7.40 Preservative: 0.1% Sodium azide Constituent: PBS
Purity	Ascites
Purification notes	Purified from ascites.
Clonality	Monoclonal
Clone number	GF5
Isotype	IgG2b

Applications

The Abpromise guarantee

Our [Abpromise guarantee](#) covers the use of ab10062 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P	★★★★★ (3)	1/100 - 1/1000. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
WB	★★★★★ (2)	Use a concentration of 1 - 5 µg/ml. Detects a band of approximately 43-45 kDa.
ELISA		Use at an assay dependent concentration.
IHC-FoFr		Use at an assay dependent concentration. PubMed: 20708681
IHC-Fr	★★★★★ (4)	1/1000.
ICC/IF	★★★★★ (8)	Use at an assay dependent concentration.
Flow Cyt	★★★★★ (1)	Use 1-2µg for 10 ⁶ cells. ab170192 - Mouse monoclonal IgG2b, is suitable for use as an isotype control with this antibody.
ICC	★★★★★ (3)	1/100.

Target

Function

GFAP, a class-III intermediate filament, is a cell-specific marker that, during the development of the central nervous system, distinguishes astrocytes from other glial cells.

Tissue specificity

Expressed in cells lacking fibronectin.

Involvement in disease

Defects in GFAP are a cause of Alexander disease (ALEXD) [MIM:203450]. Alexander disease is a rare disorder of the central nervous system. It is a progressive leukoencephalopathy whose hallmark is the widespread accumulation of Rosenthal fibers which are cytoplasmic inclusions in astrocytes. The most common form affects infants and young children, and is characterized by progressive failure of central myelination, usually leading to death usually within the first decade. Infants with Alexander disease develop a leukoencephalopathy with macrocephaly, seizures, and psychomotor retardation. Patients with juvenile or adult forms typically experience ataxia, bulbar signs and spasticity, and a more slowly progressive course.

Sequence similarities

Belongs to the intermediate filament family.


Post-translational modifications

Phosphorylated by PKN1.

Cellular localization

Cytoplasm. Associated with intermediate filaments.


Images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GFAP antibody [GF5] (ab10062)

IHC image of GFAP staining in a formalin fixed, paraffin embedded rat normal brain tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab10062 at 1/100 dilution for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.


For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



Immunocytochemistry - Anti-GFAP antibody [GF5] (ab10062)

This image is courtesy of Randal Moldrich, CNRS UMR7637, ESPCI, France


Mouse monoclonal to GFAP [GF5] (ab10062) was used in fixed murine cultures (mixed neurons/glia) at 1/100 overnight at 4°C. A secondary goat anti-mouse antibody was used for detection (Alexa Fluor 568; 1/400). Microscopy revealed diffuse cytosolic labelling. Coounterstaining with TO-PRO-3 (Molecular Probes; 660nm (converted here to blue colour) was used to identify the nucleus. The “fibrous” anti-GFAP staining of murine mixed cultures is typical of what is expected.



Flow Cytometry - Anti-GFAP antibody [GF5] (ab10062)

Overlay histogram showing SH-SY5Y cells stained with ab10062 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab10062, 1µg/1x10⁶ cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) (ab96879) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG2b [PLPV219] (ab91366, 2µg/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a positive signal in SH-SY5Y cells fixed with 4% paraformaldehyde (10


min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GFAP antibody [GF5] (ab10062)

IHC image of GFAP staining in a formalin fixed, paraffin embedded human normal hippocampus tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab10062 at 1/500 dilution for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GFAP antibody [GF5] (ab10062)

This image is courtesy of an Abreview submitted by Mr Carl Hobbs

GFAP antibody [GF5] - Astrocyte Marker


(ab10062) immunocytochemical detection in stimulated Cor1 cells. Stimulated Cor1 cells were fixed in formaldehyde, permeabilized, blocked in 1% BSA for 10 mins @ rt°C. Primary Antibody ab10062 incubated at 1/1500 for 2 hours in TBS/BSA/azide/0.3% triton. Secondary Antibody: anti mouse IgG Conjugated to: Alexa Fluor® 488 (1/1000).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GFAP antibody [GF5] (ab10062)

This image is courtesy of Carl Hobbs, King's College London, United Kingdom

GFAP antibody [GF5] - Astrocyte Marker (ab10062; 1/250 for 16h) used in Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) on Rat Tissue sections (adult brain: lateral ventricle showing astrocytes). Antigen retrieval step: Heat mediated. Blocking step: 1% BSA for 10 mins at RT. Secondary Antibody: biotin labelled goat anti mouse Ig (1/200). This image shows the exit point for the progenitor olfactory neurones, of the lateral ventricular subventricular zone.



Immunocytochemistry - Anti-GFAP antibody [GF5] (ab10062)

This image is courtesy of Randal Moldrich, CNRS UMR7637, ESPCI, France

Mouse monoclonal to GFAP [GF5] (ab10062) was used in fixed rat glial cultures at a dilution of 1/100, and incubated overnight at 4°C. Alexa Fluor 568 (1/400) secondary goat anti-mouse antibody was used for detection. Fluorescence microscopy revealed diffuse cytosolic labelling. Counterstaining with TO-PRO-3 (Molecular Probes; 660nm (converted here to blue colour) was used to identify the nucleus. The "fibrous" anti-GFAP staining of murine mixed cultures is typical of what is expected.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors