abcam

Product datasheet

Anti-Retinoic Acid Receptor alpha antibody ab28767

2 Abreviews 16 References 1 Image

Overview

Product name	Anti-Retinoic Acid Receptor alpha antibody
Description	Goat polyclonal to Retinoic Acid Receptor alpha
Host species	Goat
Specificity	This antibody is expected to recognize all three reported [human] isoforms (NP_000955.1; NP_001019980.1; NP_001028775.1).
Tested applications	**Suitable for:** WB
Species reactivity	**Reacts with:** Human **Predicted to work with:** Mouse, Chicken, Hamster, Cow, Dog
Immunogen	Synthetic peptide corresponding to Rat Retinoic Acid Receptor alpha aa 445-462 (C terminal). The immunogen is from mouse species. Sequence: C-SPSLSPSSH RSSPATQSP Database link: P11416 (Peptide available as ab4910) Run BLAST with Run BLAST with
Positive control	WB: Human breast and Human breast cancer tissue lysates.
General notes	The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing. If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	pH: 7.30 Preservative: 0.02% Sodium azide

	Constituents: 0.5% Tris buffered saline, 0.5% BSA
Purity	Immunogen affinity purified
Purification notes	This antibody was purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Clonality	Polyclonal
Isotype	IgG

Applications

The Abpromise guarantee Our Abpromise guarantee covers the use of ab28767 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
WB		Use a concentration of 0.5 - 2 µg/ml. Detects a band of approximately 55 kDa (predicted molecular weight: 51 kDa).Can be blocked with Retinoic Acid Receptor alpha peptide (ab4910). We recommend blocking in milk (3% milk in TBST) for 1 hour at room temperature.

Target

Function	Receptor for retinoic acid. Retinoic acid receptors bind as heterodimers to their target response elements in response to their ligands, all-trans or 9-cis retinoic acid, and regulate gene expression in various biological processes. The RXR/RAR heterodimers bind to the retinoic acid response elements (RARE) composed of tandem 5'-AGGTCA-3' sites known as DR1-DR5. In the absence of ligand, the RXR-RAR heterodimers associate with a multiprotein complex containing transcription corepressors that induce histone acetylation, chromatin condensation and transcriptional suppression. On ligand binding, the corepressors dissociate from the receptors and associate with the coactivators leading to transcriptional activation. RARA plays an essential role in the regulation of retinoic acid-induced germ cell development during spermatogenesis. Has a role in the survival of early spermatocytes at the beginning prophase of meiosis. In Sertoli cells, may promote the survival and development of early meiotic prophase spermatocytes. In concert with RARG, required for skeletal growth, matrix homeostasis and growth plate function (By similarity). Regulates expression of target genes in a ligand-dependent manner by recruiting chromatin complexes containing MLL5. Mediates retinoic acid-induced granulopoiesis.
Involvement in disease	Note=Chromosomal aberrations involving RARA are commonly found in acute promyelocytic leukemia. Translocation t(11;17)(q32;q21) with ZBTB16/PLZF; translocation t(15;17)(q21;q21) with PML; translocation t(5;17)(q32;q11) with NPM. The PML-RARA oncoprotein requires both the PML ring structure and coiled-coil domain for both interaction with UBE2I, nuclear microspeckle location and sumoylation. In addition, the coiled-coil domain functions in blocking RA-mediated transactivation and cell differentiation.
Sequence similarities	Belongs to the nuclear hormone receptor family. NR1 subfamily. Contains 1 nuclear receptor DNA-binding domain.
Domain	Composed of three domains: a modulating N-terminal domain, a DNA-binding domain and a C-terminal ligand-binding domain.
Post-translational	Phosphorylated on serine and threonine residues. Phosphorylation does not change during cell

modifications	cycle. Phosphorylation on Ser-77 is crucial for transcriptional activity (By similarity). Phosphorylation by AKT1 is required for the repressor activity but has no effect on DNA binding, protein stability nor subcellular localization. Phosporylated by PKA in vitro. This phosphorylation on Ser-219 and Ser-369 is critical for ligand binding, nuclear localization and transcriptional activity in response to FSH signaling. Sumoylated by SUMO2, mainly on Lys-399 which is also required for SENP6 binding. On all-trans retinoic acid (ATRA) binding, a confromational change may occur that allows sumoylation on two additional site, Lys-166 and Lys-171. Probably desumoylated by SENP6. Sumoylation levels determine nuclear localization and regulate ATRA-mediated transcriptional activity. Trimethylation enhances heterodimerization with RXRA and positively modulates the transcriptional activation. Ubiquitinated.
Cellular localization	Nucleus. Cytoplasm. Nuclear localization depends on ligand binding, phosphorylation and sumoylation. Transloaction to the nucleus in the absence of ligand is dependent on activation of PKC and the downstream MAPK phosphorylation.

Images



Western blot - Anti-Retinoic Acid Receptor alpha antibody (ab28767)

All lanes : Anti-Retinoic Acid Receptor alpha antibody (ab28767) at 1 µg/ml

Lane 1 : Human Breast tissue lysate
Lane 2 : Human Breast cancer tissue lysate

Lysates/proteins at 35 µg per lane.

Predicted band size: 51 kDa
Observed band size: 55 kDa

Primary incubation was 1 hour. Detected by chemiluminescence.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors